

Racord®

Twisted Gingival Retraction Cord with Epinephrine

Racord is twisted cotton cord saturated with a solution of racemic epinephrine hydrochloride and dried by a technique which retains the activity of the medication. Normal tissue moisture reactivates the medication, which promptly retracts gingival tissue in a manner that causes no tissue damage, yet fully exposes the gingival margin.

Contents: Each bottle contains 72 inches twisted cotton cord. The average concentration of Epinephrine for each size is:

Racord 7/00	0.30 mg/in
Racord 8/0	0.50 mg/in
Racord 9/1	0.85 mg/in
Racord 10/2	1.15 mg/in

Indications for Use: Use whenever gingival retraction is advisable in dental practice, as preceding impression taking with any elastic impression material; or prior to placement of caps, crowns, bridges; and during tooth preparation.

Contraindications: Racord is contraindicated for persons who are epinephrine sensitive, or for persons suffering from cardiovascular disease, diabetes, or hyperthyroidism. Caution should also be taken for elderly, pregnant, and pediatric patients. For such persons Pascord® is the retraction cord of choice. It is packaged in four sizes, like Racord, but contains no epinephrine.

Warnings: Contains Sodium Bisulfite, a sulfite that may cause allergic-type reactions, including anaphylactic symptoms, and life-threatening or less severe asthmatic episodes, in certain susceptible persons. The overall prevalence of sulfite sensitivity in the general population is unknown, but probably low. Sulfite sensitivity is seen more in asthmatic than non-asthmatic persons.

Precautions:

- Read and follow all instructions.
- For professional in-office use only.
- For topical intrabral use only. Do not ingest.
- Keep out of reach of children.
- Do not use on patients with Epinephrine allergy or sensitivity.

Amount to Use: Use as much cord of the appropriate sizes as is necessary to go all the way around the neck of each tooth to be retracted, with a slight overlap. Two loops may be packed around each tooth (see below), if sulcus space allows. Retraction with an epinephrine cord should

be limited to the equivalent of one normal quadrant of the mouth per appointment.

Directions for Use:

- For most effective use of Racord, loop a strand of smaller cord around the tooth and over the crevice.
- Trim the ends for a slight overlap, and pack the loop firmly into the sulcus.
- Apply a second loop of larger cord over the smaller cord in the same manner.
- When ready to apply the impression material, remove the larger cord loop only, leaving the smaller cord in place below the gingival margin until the impression procedure has been completed.
- Once the impression procedure has been completed the smaller cord is removed also.
- Alternately, a single loop of cord will suffice.
- Racord completely retracts tissue within five to eight minutes and should then be removed from the mouth.

Dispensed portions of cord are intended for single use only and should be disposed of immediately after use.

For product MSDS information

please see company website at

pascaldental.com or contact Pascal directly.

Pascal's Racord is free of gluten, soy, milk, egg, peanut, and tree nut products. Racord contains no artificial flavors or colors.

Brochures describing our latest gingival retraction materials are available upon request, or visit us at pascaldental.com.

Pascal®

International, Inc.

Rx ONLY (in USA)
For professional use only. Read
package insert for additional
cautions and warnings. This
product is not sterile. Single use
only. Dispensed portions not to
be reused.



2929 NE Northup Way
Bellevue, WA 98004 USA
425.827.4694 800.426.8051
www.pascaldental.com



OPEN
HERE

www.pascaldental.com